

algesal suractive[®] cream

Antirheumatic - analgesic - anti-inflammatory

Composition

Each 100gm contains:

Salicylic acid	6.54gm
Diethylamine	3.46gm.
Myrtecaïne	1.00gm.

Properties:

Each 100gm of algesal suractive cream contains 10gm of diethylamine salicylate.

Diethylamine salicylate has a great power of skin penetration (3 to 7 times superior to common salicylates) and dissociates in the deep inner skin tissues into diethylamine and salicylic acid.

Salicylic acid: Provides anti-rheumatic, anti-inflammatory, and analgesic action.

Diethylamine: Provides rapid analgesic action, relieving deep seated pain.

Myrtecaïne: Surface and conduction anesthetic, completes the deep analgesic and anti-inflammatory actions of diethylamine salicylate by facilitating its penetration. Also myrtecaïne has a muscle relaxant effect.

Excipient: In addition, the formula contains a special surfactant that increases and accelerates transcutaneous penetration and strengthens the analgesic action of myrtecaïne.

Indications:

• Topical treatment for fast relief of:

- Tendino -ligamentous pains.
- Muscular rheumatic pains.
- Musculo -skeletal pains.
- Inflammations

In the following conditions:

* Arthritis	* Sciatica	* Bruises
* Osteoarthritis	* Torticollis	* Synovitis
* Tendinitis	* Fibrositis	* Stiffness
* Lumbago	* Dorsalgia	* Neuralgia
* Contusions	* Low back pain	* Traumatic pains
* Disc prolapse	* Strains and sprains	* Muscular cramps

• As massage cream for sportsmen after competition, sports injuries and functional rehabilitation.

Dosage and administration

Reserved for use in adults and children over the age of 7 years.

Rub gently on the painful area, repeat massage 2 or 3 times a day.

Contraindications

- Hypersensitivity to any of the components.
- Do not apply to mucous membranes, the eyes and weeping or infected wounds.

Side effects

In very rare cases, allergic reactions may occur. If this type of reaction occurs stop taking your treatment and consult your doctor.

Pregnancy - Breast feeding: As a general rule, you should consult your doctor or pharmacist before using any drug during pregnancy or lactation.

Precaution for use: After use, it is necessary to wash your hands to prevent accidental contact between the cream and the eyes or mucous membranes.

Presentation

Tube of 40g.

Produced by:
Chemipharm Pharmaceutical Industries
6th October City - Egypt
